

Melanin-concentrating hormone directly inhibits GnRH neurons and blocks kisspeptin activation, linking energy balance to reproduction

Min Wu^a, Iryna Dumalska^a, Elena Morozova^a, Anthony van den Pol^b, and Meenakshi Alreja^{a,c,1}

Departments of ^aPsychiatry, ^bNeurosurgery, and ^cNeurobiology, Yale University School of Medicine and the Ribicoff Research Facilities, Connecticut Mental Health Center, New Haven, CT 06508

Edited by Wylie Vale, The Salk Institute for Biological Studies, La Jolla, CA, and approved August 18, 2009 (received for review July 29, 2009)

A link between energy balance and reproduction is critical for the survival of all species. Energy-consuming reproductive processes need to be aborted in the face of a negative energy balance, yet knowledge of the pathways mediating this link remains limited. Fasting and food restriction that inhibit fertility also upregulate the hypothalamic melanin-concentrating hormone (MCH) system that promotes feeding and decreases energy expenditure; MCH knockout mice are lean and have a higher metabolism but remain fertile. MCH also modulates sleep, drug abuse behavior, and mood, and MCH receptor antagonists are currently being developed as antiobesity and antidepressant drugs. Despite the clinical implications of MCH, the direct postsynaptic effects of MCH have never been reported in CNS neurons. Using patch-clamp recordings in brain slices from multiple lines of transgenic GFP mice, we demonstrate a strong inhibitory effect of MCH on an exclusive population of septal vGluT2-GnRH neurons that is activated by the puberty-triggering and preovulatory luteinizing hormone surge-mediating peptide, kisspeptin. MCH has no effect on kisspeptin-insensitive GnRH, vGluT2, cholinergic, or GABAergic neurons located within the same nucleus. The inhibitory effects of MCH are reproducible and nondesensitizing and are mediated via a direct postsynaptic Ba²⁺-sensitive K⁺ channel mechanism involving the MCHR1 receptor. MCH immunoreactive fibers are in close proximity to vGluT2-GFP and GnRH-GFP neurons. Importantly, MCH blocks the excitatory effect of kisspeptin on vGluT2-GnRH neurons. Considering the role of MCH in regulating energy balance and of GnRH and kisspeptin in triggering puberty and maintaining fertility, MCH may provide a critical link between energy balance and reproduction directly at the level of the kisspeptin-activated vGluT2-GnRH neuron.

fertility | gonadotropins | HPG axis | obesity | starvation

Nutritional status and availability of energy stores exert a profound impact on reproductive function (1–3). Reproduction is an expensive energy-consuming process, and thus it is important that puberty, pregnancy, and lactation occur only when metabolic fuel is available (4). Availability of metabolic fuel is conveyed to the brain by peripherally generated signals, such as leptin, insulin, and ghrelin, as well as by centrally released peptides such as neuropeptide Y, melanocortins, and melanin-concentrating hormone (MCH). One or more of these signals can directly or indirectly link energy balance with reproduction at one or more levels of the hypothalamic-pituitary-gonadal (HPG) axis. Thus, insulin and leptin may indirectly influence GnRH neurons (2, 5–7); leptin could regulate the HPG axis via kisspeptin-containing hypothalamic neurons (8, 9). Kisspeptin and its receptor are critical for reproduction (10–12); both kisspeptin and kisspeptin receptor knockout mice fail to enter puberty, and humans with loss of function mutations in the kisspeptin receptor exhibit hypogonadotropic hypogonadism and are infertile (13–15). Mechanistically, kisspeptin is a potent activator of GnRH neurons (16–19); it enhances pituitary gonadotropin release, triggering a cascade that is essential for entering puberty (13, 20) and for maintaining ovulation and fertility (21).

MCH neurons, which are mostly located in the lateral hypothalamus and in the zona incerta, may also target GnRH neurons directly, as MCH fibers are in close apposition with GnRH neurons (22). MCH acts via the G-protein-coupled receptors MCHR1 (23–27) and MCHR2 (28–30); only MCHR1 is present in the rodent brain, and 50–55% of rat GnRH neurons express MCHR1 (22). Intracerebral infusions of MCH can suppress (31) or enhance (32, 33) pituitary gonadotropin release, depending on the estrogenic milieu. Fasting and food restriction, which has an inhibitory effect on fertility as evidenced by decreased circulating gonadotropins and anovulation (34), upregulates the MCH system (35, 36). An activated MCH system decreases energy expenditure and increases food intake. In contrast, MCH deficiency in mice leads to leanness and increased metabolism. However, despite their leanness, MCH knockout (37, 38) and MCHR1 knockouts (39) remain fertile. The MCH system may also be involved in the regulation of sleep (40–42), drug addiction (43, 44), and mood (45). MCHR1 antagonists are currently being developed as antiobesity and antidepressant drugs (46, 47).

Despite the various functions attributed to MCH, the dramatic phenotype of the MCH knockout mice and the clinical importance of MCH, there is little information on the electrophysiological effects of MCH on CNS neurons. Available studies are restricted to hypothalamic neurons that respond to MCH with presynaptic inhibition of transmitter release (48, 49). Direct effects of MCH on membrane potential of CNS neurons have never been reported.

MCH neurons densely innervate the rodent medial septum/diagonal band of Broca (MSDB) (50, 51) that contains cholinergic, GABAergic, and vGluT2-glutamatergic neurons as well as two subpopulations of GnRH neurons, only one of which is activated by the puberty-triggering peptide kisspeptin (19).

In an effort to identify the cellular targets of MCH in the brain and to understand the mechanisms linking energy balance to reproduction, the goal of the present study was to test the hypothesis that MCH would modulate the activity of kisspeptin-activated and kisspeptin-insensitive MSDB GnRH neurons. Using electrophysiological recordings and anatomical labeling methods in multiple lines of transgenic GFP mice, we report strong, inhibitory effects of MCH on kisspeptin-activated vGluT2-GnRH neurons. Kisspeptin-insensitive GnRH, cholinergic, GABAergic, and glutamatergic MSDB neurons did not respond to MCH. The observed inhibitory effects of MCH are mediated via a direct postsynaptic mechanism and are of greater magnitude than has been reported for any CNS neuron. These actions of MCH on kisspeptin-activated GnRH

Author contributions: M.A. designed research; M.W., I.D., E.M., and A.v.d.P. performed research; A.v.d.P. contributed new reagents/analytic tools; M.W. and I.D. analyzed data; and M.A. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. E-mail: Meenakshi.Alreja@yale.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0908200106/DCSupplemental.

neurons may provide a critical link between energy balance and reproductive physiology.

Results

MCH Inhibits vGluT2-GnRH Neurons via a MCHR1-Transduced Direct Postsynaptic Mechanism. MCH inhibited a unique subpopulation of vGluT2-GnRH neurons in the MSDB that can be identified in slices prepared from either vGluT2-GFP or GnRH-GFP mice (19, 52). MCH (1 μ M) produced a 3- to 20-mV hyperpolarization (mean: 8.1 ± 1.5 mV; $n = 15$) in 59% of vGluT2-GnRH neurons recorded in brain slices prepared from 17 postpubertal mice ($n = 22$ cells; 35–160 days of age). In brain slices prepared from 300 prepubertal mice (12–30 days of age), MCH inhibited 60% of neurons tested ($n = 327$) and produced a statistically similar hyperpolarization of 6.7 ± 0.3 mV (range 2–20 mV; $n = 149$). Because neurons from both prepubertal and postpubertal mice responded similarly to MCH, the remaining studies were conducted in slices prepared from prepubertal mice.

The inhibitory effect of MCH was similar in amplitude in both prepubertal males and females and showed little desensitization to a second application of agonist administered 5–15 min after the first (females, first MCH application: 8.2 ± 0.7 mV, second MCH application: 8.1 ± 0.7 mV; $n = 20$ cells recorded from 18 mice; males, first MCH application: 7.8 ± 0.8 mV, second MCH application: 7.9 ± 0.9 mV; $n = 15$ cells recorded from 14 mice; Student's paired t test, not significant; Fig. 1*A* and *B*). Neuropeptide-glutamic acid-isoleucine ($n = 9$), cocaine- and amphetamine-regulated transcript ($n = 11$) and nesfatin ($n = 10$) had no effect on MCH responsive or nonresponsive vGluT2-GnRH neurons; all three peptides can be produced by MCH neurons (51, 53, 54). Similarly, leptin ($n = 5$) and NPY ($n = 19$), which also signal availability of energy stores, had no effect on the kisspeptin-activated vGluT2-GnRH neurons.

The MCH-induced hyperpolarization in vGluT2-GnRH neurons persisted in TTX (control: 7.1 ± 1.3 mV; TTX: 7 ± 1 mV; $n = 7$; not significant, Student's paired t test) and in “zero” Ca^{2+} /high Mg^{2+} ACSF (control: 6.5 ± 1.5 mV; “zero” Ca^{2+} /high Mg^{2+} : 8.5 ± 2 mV; $n = 8$ cells; not significant), suggesting involvement of a direct postsynaptic mechanism (Fig. 1*C–F*). As expected, the MCH-induced inhibitory effect was blocked by the MCHR1-antagonist PMC-3881-PI (control: 9.8 ± 1.4 mV; antagonist: 0 ± 0 mV; $n = 6$; $P < 0.001$, Student's paired t test; Fig. 1*G* and *H*).

To prevent the confounding effects of small changes in membrane resistance, the conductance change associated with the MCH inhibition was measured under voltage-clamp conditions. At a holding potential of -65 mV, MCH produced a 6.5 ± 1 pA outward current (Fig. 1*I* and *K*) that was associated with a significant increase in membrane conductance (control: 0.8 ± 0.2 nS; MCH: 1 ± 0.2 nS; $P < 0.01$; $n = 22$), suggesting a net opening of channels. The MCH-induced outward current was significantly reduced in the presence of BaCl_2 , a blocker of K^+ channels (control: 11.6 ± 1.4 pA; Ba^{2+} : 1.8 ± 0.4 pA; $n = 5$; Student's paired t test; $P < 0.01$; Fig. 1*I* and *J*). Ba^{2+} alone produced a 10.6 ± 3.3 pA inward current in voltage-clamp recordings ($n = 5$). Consistent with the involvement of a K^+ current, MCH-induced outward current reversed at mean reversal potential of -101 ± 1.7 mV ($n = 5$), which is near the calculated E_K of -101 mV (Fig. 1*L*). Thus, MCH-induced inhibition of vGluT2-GnRH neurons involves opening of Ba^{2+} -sensitive K^+ channels.

MCH Selectively Inhibits a Subpopulation of Kisspeptin-Activated vGluT2-GnRH Neurons in the MSDB. The above-described effects of MCH occurred in a very unique population of neurons within the MSDB. The MCH-inhibited vGluT2-GnRH neurons could be distinguished from other major neuronal populations within the MSDB by their lack of excitatory response to the Group I glutamate metabotropic-receptor agonist DHPG and by their strong and persistent activation by the neuropeptide, kisspeptin,

the natural ligand of GPR54 (19, 52). Although DHPG-sensitivity was examined in every neuron tested, because of the strong and prolonged nature of the kisspeptin response, kisspeptin agonist was applied only at the end of the experiment. Sensitivity to kisspeptin was assessed using the bioactive fragment kisspeptin-10 (KP-10). A total of 75 MCH-responsive vGluT2-GnRH neurons were confirmed to be KP-10-sensitive at the end of the experiment (Fig. 2*A*, *B*, and *E*).

MCH had no effect on the 36 KP-10-insensitive GnRH-GFP cells that were recorded in brain slices prepared from 30 mice (Fig. 2*C* and *E*) or on the 47 KP-10-insensitive vGluT2-GFP neurons that were recorded from 42 mice (Fig. 2*D* and *E*); these neurons were strongly activated by DHPG. MCH also had no effect on the cholinergic ($n = 27$) or on the GABAergic neurons ($n = 12$; Fig. 2*E*); both neuronal populations are similarly activated by DHPG but not by KP-10 (19, 55, 56). Thus, MCH selectively inhibits the KP-10-activated, DHPG-insensitive neurons within the MSDB.

MCH Interrupts Kisspeptin-Induced Activation of vGluT2-GnRH Neurons.

Because of a strong link between energy balance and reproduction, we next determined whether MCH, by virtue of its inhibitory activity, could oppose the long-lasting excitatory effect of kisspeptin. Under control conditions, KP-10 activation lasts an average of 16 ± 1.5 min (Fig. 3*A*) (19). In 12 cells, 100 nM KP-10 was applied for 5 s and, after establishment of the excitatory response to KP-10, MCH was applied for 15 s. MCH interrupted the excitatory effect of KP-10 and produced a 10.9 ± 1.1 mV hyperpolarization (Fig. 3*B*). This interruption lasted an average of 1.8 ± 0.3 min ($n = 12$). In addition, because of the nondesensitizing nature of the MCH response, repeated applications of MCH continued to block KP-10 activation (Fig. 3*C*). A similar interruption was observed in voltage-clamp recordings ($n = 3$) (Fig. 3*D*).

MCH Immunoreactive Fibers Are Present in the Vicinity of Septal vGluT2-GnRH Neurons.

Because MCH inhibited KP-10-activated vGluT2-GnRH neurons, we specifically determined the anatomical relationship between MCH-immunoreactive fibers and vGluT2-GFP and GnRH-GFP neurons using vGluT2-GFP ($n = 5$) and GnRH-GFP ($n = 5$) mice and a well-characterized antisera against MCH (51). In sections prepared from vGluT2-GFP mice, MCH immunoreactive axons stained red were found throughout the MSDB and appeared in close juxtaposition to vGluT2 cell bodies and dendrites. In some cells, many MCH boutons appeared to contact single GFP cells (Fig. 4*A–D*). Similarly, in the GnRH-GFP mice, MCH axons appeared in close juxtaposition to GFP cell bodies and dendrites (Fig. 4*E–G*). Elimination of either the primary or secondary antibody gave no staining, and substitution with antisera against hypocretin/orexin resulted in a different pattern of staining. Thus MCH axons contact GnRH as well as the vGluT2 neurons. Ultrastructural analysis would help to address the question of whether these contacts are synaptic in nature.

Discussion

In the present study, we report a direct postsynaptic inhibitory action of the orexigenic peptide MCH on CNS neurons for the first time. These effects suggest an additional function of MCH in the brain. The inhibitory effects of MCH occur in an exclusive subset of basal forebrain neurons, namely, the vGluT2-GnRH neurons. The MCH-induced inhibition in vGluT2-GnRH neurons is mediated via a MCHR1-receptor-transduced postsynaptic mechanism that involves an opening of Ba^{2+} -sensitive K^+ channels. MCH-immunoreactive fibers are located in close apposition to vGluT2-GFP and GnRH-GFP neurons. Interestingly, the inhibitory effects of MCH can interrupt or block the persistent excitatory effect of the hypothalamic peptide kisspeptin, which is essential for reproduction. Given the role of MCH in energy balance and of kisspeptin in triggering puberty and in sustaining ovulation and fertility, MCH inhibition of kisspeptin-sensitive vGluT2-GnRH neurons suggests a




Fig. 1. MCH inhibits vGluT2-GnRH neurons via a direct postsynaptic mechanism involving MCH1R and an opening of Ba^{2+} -sensitive K^+ channels. (A) Chart record from an example current-clamp recording shows that two consecutive applications of 1 μ M MCH at an interval of 8.5 min produced a comparable hyperpolarization of 13 mV. (B) Bar charts summarize the data for two consecutive applications of MCH applied at intervals of 5–15 min. Note the similar magnitude of the response in the two sexes, the reproducibility of the inhibitory effects, and a lack of desensitization. (C–F) Chart records and bar charts show that MCH inhibition does not change in the presence of TTX or in “zero” Ca^{2+} /high Mg^{2+} ACSF. (G, H) Chart record and bar chart shows the

MCH-mediated link between energy balance and reproduction directly at the level of the GnRH neuron.


In the present study, brief 15-s applications of MCH produced a strong and reproducible inhibition in a highly selective population of vGluT2-GnRH neurons that is persistently activated by the Gq-coupled receptor GPR54 ligand kisspeptin but not by the Gq-coupled Group I glutamate metabotropic receptor agonist DHPG. In contrast, MCH had no effect on DHPG-activated, kisspeptin-insensitive GnRH, vGluT2, cholinergic or GABAergic neurons that are located within the same basal forebrain area. Thus the observed inhibitory effects of MCH in the MSDB are highly selective for a neuronal population that is involved in triggering puberty and sustaining key reproductive functions that are essential to fertility. Therefore, these findings suggest a novel function of MCH. In addition, the absence of MCH inhibition in kisspeptin-insensitive, DHPG-activated GnRH neurons provides an additional line of evidence in favor of two populations of GnRH neurons that serve different functions by virtue of their neuro-modulator responses (19).

MCH has been shown to alter the firing rate of ventromedial hypothalamic and arcuate neurons; whether these effects are mediated via a direct mechanism has not been determined (57). In cultured lateral hypothalamic neurons, MCH inhibits voltage-dependent calcium channels and decreases both glutamatergic and GABAergic synaptic transmission without involvement of K^+ currents and without having a direct effect on membrane potential and conductance (48, 49). In the present study, conducted in murine brain slices, MCH produced a 2- to 20-mV hyperpolarization and a marked increase in membrane conductance that persisted after blockade of synaptic transmission, suggesting a direct effect of MCH on membrane properties. These effects suggest an additional function of MCH in the brain. The MCH-induced changes in membrane properties reversed near E_{K} and did not occur in the presence of barium, suggesting involvement of K^+ channels. Considering that MCH1R is coupled to the G proteins, Gi and Go, it is not surprising that MCH is inhibitory. Nevertheless, earlier studies in non-neuronal kidney cell lines found that MCH raised intracellular calcium (25, 26). In neurons, an increase in calcium is generally associated with excitation, underlining the complications of studying neuronal receptors in non-neurons.

Unlike MCH, NEI, CART, or nesfatin had no apparent effect on the membrane properties of kisspeptin-activated vGluT2-GnRH neurons. The absence of leptin and NPY effects on these neurons is consistent with the absence of leptin receptors in GnRH neurons (6); NPY effects have been reported only in lactating mice (58).

The physiological significance of the observed inhibitory effects of MCH is underscored by our MCH immunoreactivity studies. The high density of MCH fibers that we observed in MSDB in mice is consistent with earlier reports in rats (50, 51, 59). The close appositions observed between GnRH-GFP neurons and MCH-ir fibers in mice are consistent with the study in rats (22). In addition, in our study, close appositions were noted between vGluT2-GFP neurons and MCH fibers that are consistent with the observed inhibitory effects of MCH on vGluT2-GnRH neurons.

effect of MCH before and after bath application of a MCH1R antagonist. Broken lines indicate base line, and number in mV denotes resting membrane potential in quiescent neurons. Downward vertical deflections are in response to -0.01 nA current pulses delivered every 4 s to monitor the input resistance of the recorded neuron. (I, J) In an example voltage-clamp recording (holding potential: -65 mV) MCH produced a 15-pA outward current. Bath-applied BaCl_2 , a K^+ channel blocker, produced a 6-pA inward current and blocked MCH inhibition. (K) Cell in which MCH induced a 7-pA outward current that is closer to the mean current observed under our experimental conditions. (L) I-V curves, obtained using slow steady-state ramps in a voltage-clamped neuron, in the absence and in the presence of MCH. Subtracted current is also shown. The MCH current reversed at -103 mV, close to the calculated E_{K} of -101 mV.



PNAS | October 6, 2009 | vol. 106 | no. 40 | 17221

GFP mice. MCH and other agonists were applied using a Y-tube. Immunocytochemistry was performed to determine MCH innervation of vGluT2-GnRH neurons using a well-established antibody [for detailed methods, see [S1](#)].

ACKNOWLEDGMENTS. This work was supported by National Institution of Health grants MH61465, NS41454, and NS48476, and by the State of Connecticut, Department of Mental Health and Addiction Services.

- Schneider JE (2004) Energy balance and reproduction. *Physiol Behav* 81:289–317.
- Crown A, Clifton DK, Steiner RA (2007) Neuropeptide signaling in the integration of metabolism and reproduction. *Neuroendocrinology* 86:175–182.
- Hill JW, Elmquist JK, Elias CF (2008) Hypothalamic pathways linking energy balance and reproduction. *Am J Physiol Endocrinol Metab* 294:E827–E832.
- Mircea CN, Lujan ME, Pierson RA (2007) Metabolic fuel and clinical implications for female reproduction. *J Obstet Gynaecol Can* 29:887–902.
- Gottsch ML, Clifton DK, Steiner RA (2004) Galanin-like peptide as a link in the integration of metabolism and reproduction. *Trends Endocrinol Metab* 15:215–221.
- Quenell JH, et al. (2009) Leptin indirectly regulates GnRH neuronal function. *Endocrinology*.
- Leshan RL, et al. (2009) Direct innervation of GnRH neurons by metabolic- and sexual odorant-sensing leptin receptor neurons in the hypothalamic ventral premammillary nucleus. *J Neurosci* 29:3138–3147.
- Smith JT, Acohido BV, Clifton DK, Steiner RA (2006) Kiss-1 neurones are direct targets for leptin in the ob/ob mouse. *J Neuroendocrinol* 18:298–303.
- Castellano JM, et al. (2006) Expression of hypothalamic Kiss-1 system and rescue of defective gonadotropic responses by kisspeptin in streptozotocin-induced diabetic male rats. *Diabetes* 55:2602–2610.
- Kauffman AS, Clifton DK, Steiner RA (2007) Emerging ideas about kisspeptin- GPR54 signaling in the neuroendocrine regulation of reproduction. *Trends Neurosci* 30:504–511.
- Chan YM, Broder-Fingert S, Seminara SB (2009) Reproductive functions of kisspeptin and Gpr54 across the life cycle of mice and men. *Peptides* 30:42–48.
- Gottsch ML, Clifton DK, Steiner RA (2009) From Kiss1 to kisspeptins: An historical perspective and suggested nomenclature. *Peptides* 30:4–9.
- Seminara SB, et al. (2003) The GPR54 gene as a regulator of puberty. *N Engl J Med* 349:1614–1627.
- Lapatto R, et al. (2007) Kiss1^{−/−} mice exhibit more variable hypogonadism than Gpr54^{−/−} mice. *Endocrinology* 148:4927–4936.
- d'Anglemont de Tassigny X, et al. (2007) Hypogonadotropic hypogonadism in mice lacking a functional Kiss1 gene. *Proc Natl Acad Sci USA* 104:10714–10719.
- Han SK, et al. (2005) Activation of gonadotropin-releasing hormone neurons by kisspeptin as a neuroendocrine switch for the onset of puberty. *J Neurosci* 25:11349–11356.
- Pielecka-Fortuna J, Chu Z, Moenter SM (2008) Kisspeptin acts directly and indirectly to increase gonadotropin-releasing hormone neuron activity and its effects are modulated by estradiol. *Endocrinology* 149:1979–1986.
- Zhang C, Roepke TA, Kelly MJ, Ronnekleiv OK (2008) Kisspeptin depolarizes gonadotropin-releasing hormone neurons through activation of TRPC-like cationic channels. *J Neurosci* 28:4423–4434.
- Dumalska I, et al. (2008) Excitatory effects of the puberty-initiating peptide kisspeptin and group I metabotropic glutamate receptor agonists differentiate two distinct subpopulations of gonadotropin-releasing hormone neurons. *J Neurosci* 28:8003–8013.
- de Roux N, et al. (2003) Hypogonadotropic hypogonadism due to loss of function of the Kiss1-derived peptide receptor GPR54. *Proc Natl Acad Sci USA* 100:10972–10976.
- Clarkson J, d'Anglemont de Tassigny X, Moreno AS, Colledge WH, Herbison AE (2008) Kisspeptin-GPR54 signaling is essential for preovulatory gonadotropin-releasing hormone neuron activation and the luteinizing hormone surge. *J Neurosci* 28:8691–8697.
- Williamson-Hughes PS, Grove KL, Smith MS (2005) Melanin concentrating hormone (MCH): A novel neural pathway for regulation of GnRH neurons. *Brain Res* 1041:117–124.
- Shimomura Y, et al. (1999) Isolation and identification of melanin-concentrating hormone as the endogenous ligand of the SLC-1 receptor. *Biochem Biophys Res Commun* 261:622–626.
- Bächner D, Kreienkamp H, Weise C, Buck F, Richter D (1999) Identification of melanin concentrating hormone (MCH) as the natural ligand for the orphan somatostatin-like receptor 1 (SLC-1). *FEBS Lett* 457:522–524.
- Lembo PM, et al. (1999) The receptor for the orexigenic peptide melanin-concentrating hormone is a G-protein-coupled receptor. *Nat Cell Biol* 1:267–271.
- Chambers J, et al. (1999) Melanin-concentrating hormone is the cognate ligand for the orphan G-protein-coupled receptor SLC-1. *Nature* 400:261–265.
- Saito Y, et al. (1999) Molecular characterization of the melanin-concentrating-hormone receptor. *Nature* 400:265–269.
- Wang S, et al. (2001) Identification and pharmacological characterization of a novel human melanin-concentrating hormone receptor, mch-r2. *J Biol Chem* 276:34664–34670.
- Sailer AW, et al. (2001) Identification and characterization of a second melanin-concentrating hormone receptor, MCH-2R. *Proc Natl Acad Sci USA* 98:7564–7569.
- An S, et al. (2001) Identification and characterization of a melanin-concentrating hormone receptor. *Proc Natl Acad Sci USA* 98:7576–7581.
- Tsakamura H, et al. (2000) Intracerebroventricular administration of melanin-concentrating hormone suppresses pulsatile luteinizing hormone release in the female rat. *J Neuroendocrinol* 12:529–534.
- Gonzalez MI, Baker BI, Wilson CA (1997) Stimulatory effect of melanin-concentrating hormone on luteinizing hormone release. *Neuroendocrinology* 66:254–262.
- Murray JF, et al. (2006) Evidence for a stimulatory action of melanin-concentrating hormone on luteinizing hormone release involving MCH1 and melanocortin-5 receptors. *J Neuroendocrinol* 18:157–167.
- Wade GN, Jones JE (2004) Neuroendocrinology of nutritional infertility. *Am J Physiol Regul Integr Comp Physiol* 287:R1277–R1296.
- Qu D, et al. (1996) A role for melanin-concentrating hormone in the central regulation of feeding behaviour. *Nature* 380:243–247.
- Presse F, Sorokovsky I, Max JP, Nicolaidis S, Nahon JL (1996) Melanin-concentrating hormone is a potent anorectic peptide regulated by food-deprivation and glucopenia in the rat. *Neuroscience* 71:735–745.
- Shimada M, Tritos NA, Lowell BB, Flier JS, Maratos-Flier E (1998) Mice lacking melanin-concentrating hormone are hypophagic and lean. *Nature* 396:670–674.
- Kokkotou E, et al. (2005) Mice with MCH ablation resist diet-induced obesity through strain-specific mechanisms. *Am J Physiol Regul Integr Comp Physiol* 289:R117–R124.
- Marsh DJ, et al. (2002) Melanin-concentrating hormone 1 receptor-deficient mice are lean, hyperactive, and hyperphagic and have altered metabolism. *Proc Natl Acad Sci USA* 99:3240–3245.
- Verret L, et al. (2003) A role of melanin-concentrating hormone producing neurons in the central regulation of paradoxical sleep. *BMC Neurosci* 4:19.
- Willie JT, Sinton CM, Maratos-Flier E, Yanagisawa M (2008) Abnormal response of melanin-concentrating hormone deficient mice to fasting: Hyperactivity and rapid eye movement sleep suppression. *Neuroscience* 156:819–829.
- Hassani OK, Lee MG, Jones BE (2009) Melanin-concentrating hormone neurons discharge in a reciprocal manner to orexin neurons across the sleep-wake cycle. *Proc Natl Acad Sci USA* 106:2418–2422.
- Georgescu D, et al. (2005) The hypothalamic neuropeptide melanin-concentrating hormone acts in the nucleus accumbens to modulate feeding behavior and forced-swim performance. *J Neurosci* 25:2933–2940.
- Pissios P, et al. (2008) Dysregulation of the mesolimbic dopamine system and reward in MCH^{−/−} mice. *Biol Psychiatry* 64:184–191.
- Adamantidis A, de Lecea L (2008) Sleep and metabolism: Shared circuits, new connections. *Trends Endocrinol Metab* 19:362–370.
- Nahon JL (2006) The melanocortins and melanin-concentrating hormone in the central regulation of feeding behavior and energy homeostasis. *C R Biol* 329:623–638; discussion 653–655.
- Shimazaki T, Yoshimizu T, Chaki S (2006) Melanin-concentrating hormone MCH1 receptor antagonists: A potential new approach to the treatment of depression and anxiety disorders. *CNS Drugs* 20:801–811.
- Gao XB, van den Pol AN (2002) Melanin-concentrating hormone depresses L-, N-, and P/Q-type voltage-dependent calcium channels in rat lateral hypothalamic neurons. *J Physiol* 542:273–286.
- Gao XB, van den Pol AN (2001) Melanin concentrating hormone depresses synaptic activity of glutamate and GABA neurons from rat lateral hypothalamus. *J Physiol* 533:237–252.
- Bittencourt JC, Elias CF (1998) Melanin-concentrating hormone and neuropeptide EI projections from the lateral hypothalamic area and zona incerta to the medial septal nucleus and spinal cord: A study using multiple neuronal tracers. *Brain Res* 805:1–19.
- Bittencourt JC, et al. (1992) The melanin-concentrating hormone system of the rat brain: An immuno- and hybridization histochemical characterization. *J Comp Neurol* 319:218–245.
- Wu M, Dumalska I, Morozova E, van den Pol AN, Alreja M (2009) Gonadotropin inhibitory hormone inhibits basal forebrain vGluT2-gonadotropin-releasing hormone neurons via a direct postsynaptic mechanism. *J Physiol* 587:1401–1411.
- Cvetkovic V, et al. (2004) Characterization of subpopulations of neurons producing melanin-concentrating hormone in the rat ventral diencephalon. *J Neurochem* 91:911–919.
- Fort P, et al. (2008) The satiety molecule nesfatin-1 is co-expressed with melanin concentrating hormone in tubular hypothalamic neurons of the rat. *Neuroscience* 155:174–181.
- Wu M, Hajszan T, Leranah C, Alreja M (2003) Nicotine recruits a local glutamatergic circuit to excite septohippocampal GABAergic neurons. *Eur J Neurosci* 18:1155–1168.
- Wu M, Hajszan T, Xu C, Leranah C, Alreja M (2004) Group I metabotropic glutamate receptor activation produces a direct excitation of identified septohippocampal cholinergic neurons. *J Neurophysiol* 92:1216–1225.
- Davidova H, Li Y, Plagemann A (2002) Hypothalamic ventromedial and arcuate neurons of normal and postnatally overnourished rats differ in their responses to melanin-concentrating hormone. *Regul Pept* 108:103–111.
- Xu J, Kirigiti MA, Cowley MA, Grove KL, Smith MS (2009) Suppression of basal spontaneous gonadotropin-releasing hormone neuronal activity during lactation: Role of inhibitory effects of neuropeptide Y. *Endocrinology* 150:333–340.
- Saito Y, Cheng M, Leslie FM, Civelli O (2001) Expression of the melanin-concentrating hormone (MCH) receptor mRNA in the rat brain. *J Comp Neurol* 435:26–40.
- McBriar MD (2006) Recent advances in the discovery of melanin-concentrating hormone receptor antagonists. *Curr Opin Drug Discov Devel* 9:496–508.
- Gibson MJ, Wu TJ, Miller GM, Silverman AJ (1997) What nature's knockout teaches us about GnRH activity: Hypogonadal mice and neuronal grafts. *Horm Behav* 31:212–220.
- Gallardo MG, Chiochio SR, Tramezzani JH (2004) Changes of melanin-concentrating hormone related to LHRH release in the median eminence of rats. *Brain Res* 1030:152–158.
- Ahima RS, et al. (1996) Role of leptin in the neuroendocrine response to fasting. *Nature* 382:250–252.
- Polak P, et al. (2008) Adipose-specific knockout of raptor results in lean mice with enhanced mitochondrial respiration. *Cell Metab* 8:399–410.
- Xu ZP, Wawrousek EF, Piatigorsky J (2002) Transketolase haploinsufficiency reduces adipose tissue and female fertility in mice. *Mol Cell Biol* 22:6142–6147.
- Yamada M, et al. (2001) Mice lacking the M3 muscarinic acetylcholine receptor are hypophagic and lean. *Nature* 410:207–212.
- Fischer J, et al. (2009) Inactivation of the Fto gene protects from obesity. *Nature* 458:894–898.
- Peder M, Porkka-Heiskanen T, Laakso ML, Johansson G (1989) Rapid eye movement sleep deprivation depresses plasma FSH and LH in castrated rats. *Physiol Behav* 45:1167–1170.
- Fehm HL, Clausen J, Kern W, Pietrowsky R, Born J (1991) Sleep-associated augmentation and synchronization of luteinizing hormone pulses in adult men. *Neuroendocrinology* 54:192–195.